

TITLE OF THE INVENTION β -1.3-1.6 GLUCAN (*AUREOBASIDIUM* MEDIUM)BACKGROUND OF THE INVENTIONField of the Invention

5 The invention relates to a particular type of β -glucan--a microbial polysaccharide composed of D-glucose in either straight or branched chains with glycosidic linkages. More specifically, the present invention is directed to industrial and commercial applications of a β -1.3-1.6-glucan appearing in the medium of the black yeast, *Aureobasidium*. *Aureobasidium* is one of several genera of black yeasts, characterized by mostly slow-growing, black, pasty colonies. β -1.3-1.6-glucans are complex acidic polysaccharides having a main chain comprising subunits having 1,3 linkages and branches at the 3- and 6-positions. The β -glucans of the present invention have a variety of industrial and commercial uses, including applications in pharmaceutical or medical products or treatments, for the removal or control of environmental or microbiological contaminants, in cosmetics, and in nutritional products and foods.

Description of the Related Art

10
15
20 Various microbial and fungal polysaccharides with different structures and diverse physiological properties are known. Such polysaccharides have been studied from a variety of different aspects, see e.g. Hamada Nobutake: Kobunshi Kako 9-16 (1987); and Ohno Naohito: J. Bacteria Japan 527 (2000).

25 Microbial polysaccharides are formed from different types of monosaccharide units, such as particular types of sugars, including glucose (dextrose, grape sugar) and fructose (fruit sugar). Glucans comprise units of glucose. However, depending on the type of linkage between these subunits, a glucan may be either an α -glucan or β -glucan, see Hamada, N, Tsujisaka Y: Agric. Biol. Chem. 30: 266 (1966).

 Microbial polysaccharides may have unique physical properties and physiological activities. For instance, their physical properties can provide consistency, body and

substance to foods. They can be utilized as thickening agents and as raw materials in semi-solid foods, such as jellies. Polysaccharides, such as curdlan, are utilized as antioxidant films, Szezesniak et al., J. Food Sci. 27: 381 (1962), Smiey, Food Technol 20: 112 (1966). Moreover, microbial polysaccharides which impart stability, water retention and
5 dispensability may be used in a variety of products, including cosmetics, paints, fertilizers, papermaking, fibers, etc., Jeans, Extracellular Microbiol. Polysaccharides, Sanford et al. (eds), American Chemical Society, Washington, D.C. (1977).

On the other hand, other types of polysaccharides, such as dextran, are used for certain pharmaceutical or medical applications. For instance, dextran is utilized as a plasma
10 substitute or blood flow improver. Polysaccharides, such as chizophyllan, scleroglucan, and curdlan have physiological activities that may be useful in treating diseases such as cancer, Komatsu et al., Gann 60: 137 (1969); Singh et al., Carbohydr. Res. 37: 245 (1974). In general, such polysaccharides do not act directly on cancer cells, but by enhancing immunity. Protein-containing polysaccharides of *Trametes versicolor* PSK, the polysaccharide lentinan of *Lentinus edodes* and the polysaccharide schizophyllan of *Schizophyllum commune* are used as immunotherapeutic drugs for cancer.

An acidic polysaccharide produced by strain FERM P-14228 is described by U.S. Patent 5,789,579. This patent is directed to α -glucans having α -1.4 bonding, unlike the β -glucans of the present invention that have β -1.3-1.6 linkages.

β -glucans are found in mushrooms, such as *Agaricus* and *Lentinus*, various starches, cellulose, seaweeds, red algae, yeasts and other similar micro-organisms. Some types of β -glucans play a significant role in health maintenance, see Fujii Noboru: A miracle of β -glucan, Society for Study of Beta Glucan. The present inventors have found that β -1.3-1.6glucan (*Aureobasidium medium*) contains much higher quantities of β -glucan than
25 mushrooms--over ten-times more.

Physiological Effects of β -glucans. Many studies have been made with respect to the physiological effects of β -glucans. In particular, the regulatory actions of these compounds on the vital balance and vital immune reaction have been studied. For instance, the protective effects of these compounds in cancer and infectious disease have been studied.
30 Some types of β -glucans, which are of a different type from those of the present invention,

have antitumor activities. For instance, these have been observed on 3LL solid tumor cells derived with methylcholanthrene-I in experimental animals, see Fujii Noboru, Shinohara Satoru: Report of the Department of Agriculture, Miyazaki University 33: 243 (1986).

Glucan Receptor. β -glucan receptors are known to be expressed on immunocompetent cells, such as monocytes and macrophages, see Czop et al., J. Exp Med 173: 1511 (1991), Xia et al., J. Immunol. 162: 7285 (1999), Czop, et al., Prog. Clin. Biol. Res. 297: 287 (1989). Certain β -glucans are known to induce the production of inflammatory cytokines from immunocompetent cells, for example, the interleukin 1 receptor antagonist (IL-1Ra) which is an antagonist of interleukin 1 β (IL-1 β), see Poutsika et al., Blood 82: 3695 (1993). It also induces the production of tumor necrosis factor- α (TNF- α), interleukin 1 β (IL-1 β) and platelet activating factor (PAF), and the like, see Abel et al., Int. J. Immunopharmacol. 14: 1363 (1992), Elstad, et al., J. Immunol. 52: 220 (1994).

The molecular structure of the β -glucan receptor has been found to correspond to complement receptor 3 ("CR3") that binds complement component C3. Recently, this receptor has been shown to be an integrin type CD11b/CD18 molecule itself, see Ress, et al., Complement 4: 61 (1987), Thornton, et al., J. Immunol. 156: 235 (1996). It has also been found that this CR3 (CD11b/CD18) molecule is a receptor mainly manifested on the cell surface of monocytes and macrophages and in NK cells. NK cells activated by β -glucan bonding to the CD11b/CD18 molecule via this receptor (CR3) molecule are involved in the destruction and removal of cancer cells, see Di Renzo et al., Eur. J. Immunol. 21: 1755 (1991), Ross, et al., Immunopharmacology 42: 61 (1999), Yan, et al., J. Immunol. 163: 3045 (1999).

Signal transmission. The analysis of signal transmission modes from the β -glucan receptor has been studied. $Nf_{\kappa}B$ plays a role as an intranuclear transmission substance of immuno-B-cyte which takes part closely in this signal transmission system, Wakshull et al., Immunopharmacology 41: 89 (1999). Moreover, in model experiments for the therapy of cancer, from the fact that β -glucan is a receptor (CD11b/CD18 molecule) of C3 component of complement, see Yan, et al., J Immunol 163: 3045 (1999), it is reported that the route via this receptor is effective for the destruction of cancer cells opsonized with specific antibody of cancer, see Xia et al., J. Immunol. 162: 2281 (1999).

Drug effects on glucan receptor. On the other hand, functional experiments between drugs having an influence on the immune reaction and β -glucan receptor are also being conducted, and new roles of β -glucan in the vital immune system have become clear in such ways that, in particular, β -glucan receptor (CD11b/CD18 molecule) accelerates its receptor function with glucocorticoid, and the like, see Yashioka et al., FEMS Immunol. Med. Microbiol. 21: 171 (1998), Kay, et al., Immunology 81: 96 (1994).

Glucan effects *in vivo*. The effects of β -glucans *in vivo* have also been investigated. Administration of a glucan increases the survival and longevity (macrobiotic effect) of mice exposed to lethal levels of radiation. This effect is attributed to hemopoietic effects of β -glucan and direction of immunocompetent cells into a maturation process, see Patchen et al., Methods Find. Exp. Clin. Pharmacol. 8: 151 (1986). In addition, β -glucans stimulate macrophage function and may thus provide protective effects against certain infectious diseases, such as tuberculosis, see Hetland et al., Scand. J. Immunol 47: 548 (1998). Immunization with the glucan-binding domain of *Streptococcus mutans* glucosyltransferase provides immunity against infection with *Streptococcus mutans*, see Jespersgaard et al., Infect. Immun. 67: 6543 (1999).

Immune enhancement with glucans and other substances. While some β -glucans may enhance certain components of the immune system, such as macrophage function, immunological enhancement is a complex phenomenon and at the cell level many intracellular transmission substances are involved in producing these effects. For instance, recently it has been found that mitogen activating protein kinase (MAPK) and calcium controlling the activation and management of immune system are closely involved in these phenomena, see McLeish, et al., J. Leukoc. Biol. 64: 835 (1998), Mork, et al., Immunopharmacology 40: 77 (1998).

β -glucan analogs and diagnosis. Clinically, β -glucan analogues in plasma in patients with pneumonia have been observed. For instance, β -glucan analogues increase in plasma in cases of *Pneumocystis carinii* pneumonia having no overt AIDS-like symptoms, see Teramoto et al., J. Med. Microbiol. 49: 393 (2000).

While some β -glucans were known to stimulate particular components of the cellular immune system, such as macrophage functions, the present inventors have found that the β -1.3-1.6glucan (*Aureobasidium* medium) of the present invention exerts other specific

functional activities, such as inducing increases in the numbers of lymphocytes, inducing DNA synthesis and cell division in lymphocytes, exerting direct effects on cancer cells, and directly inhibiting the proliferation of microbes.

SUMMARY OF THE INVENTION

5 The invention relates to an *Aureobasidium* medium containing β -1.3-1.6glucan as a major component and a method of producing the same by culturing fungi that produce β -1.3-1.6glucan of *Aureobasidium*. The unique functional properties of the β -glucans of the present invention, which include particular modulatory and regulatory effects on the immune system, make these products particularly advantageous for a variety of industrial and commercial applications. The glucans of the present invention may be used as present in *Aureobasidium* culture medium, which is preferably sterilized or filtered, or may be used after further purification. Such applications include use in pharmaceutical or medical products or treatments, in diagnostic products and methods, for the removal or control of environmental or microbiological contaminants, in chemical products, in cosmetics, and in nutritional products and foods.

10 The present inventors have discovered that β -1.3-1.6glucan (*Aureobasidium* medium) prepared according to the invention has unique physical properties and physiological activity quite different from those of other types of β -glucans. In particular, the β -1.3-1.6glucan (*Aureobasidium* medium) of the present invention modulates and regulates the immune system. It has been found to have a specific effect on the production of interleukin-6 (IL-6) which is involved in controlling the number of lymphocytes in human peripheral blood, in cell division and in other aspects of immunity.

15 Moreover, the β -1.3-1.6glucan (*Aureobasidium* medium) of the present invention effectively inhibited the proliferation of cancer cells, such as leukemia cells. While not being limited to a particular mechanism of action, it is considered that this effect may involve the programmed death of cancer (leukemia) cells or apoptosis. Thus, the β -1.3-1.6glucan (*Aureobasidium* medium) of the present invention has advantageous applications for clinical therapy of cancer.

20 It has also been found that β -1.3-1.6glucan (*Aureobasidium* medium) influences the

expression or function of adhesion molecules involved in immunity. For instance, the inhibition of expression or down-regulation of integrin-type adhesion molecules, such as CD11a and CD49d, expressed in monocytic cells was recognized. This finding provides for application of β -1.3-1.6glucan (*Aureobasidium* medium) in the treatment of various microbial diseases. For instance, the CD11a molecule is a receptor for rotavirus, which causes infant and child diarrhea. Since β -1.3-1.6glucan (*Aureobasidium* medium) down-regulates the expression or function of CD11a molecules as shown in the example below, it may be used to prevent the attachment or adsorption of rotavirus, and prevent or ameliorate infections involving CD11a.

Moreover, the CD49d molecule is also an integrin type adhesion molecule and an immune function regulator. This molecule is considered to be strongly involved in the infiltration of T lymphocytes into the articular synovial membrane of patients having chronic articular rheumatism. Since the expression of CD49d molecule is also subject to the down-regulation with β -1.3-1.6glucan (*Aureobasidium* medium), it may be used in methods for therapy of diseases such as chronic articular rheumatism and other autoimmune diseases, involving expression of CD49d. Thus, the β -1.3-1.6glucan (*Aureobasidium* medium) may exert anti-inflammatory effects.

Applicants have also discovered that apart from the advantageous immunological effects of β -1.3-1.6glucan (*Aureobasidium* medium), that this substance may be used for preventing the propagation of environmental microbes. Thus, β -1.3-1.6glucan (*Aureobasidium* medium) may be used as a novel antiseptic, preservative or antimicrobial agent. Such agents have application for maintaining personal cleanliness, for instance of the body, hands or feet, and as a portable disinfectant. Such a disinfectant may be used for persons who cannot bathe, such as the aged or infirm and for prevention of nosocomial infections. It may also be used to remove environmental pollutants.

As described and shown below, the present invention involving β -1.3-1.6glucan (*Aureobasidium* medium) may be advantageously and safely used in medical and pharmaceutical products and methods, in foods and cosmetics and as an antiseptic or antimicrobial agent.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the numbers of lymphocytes in peripheral blood.

Fig. 2, panels (a) and (b) compare the morphological changes in lymphocytes grown in control medium with those grown in medium containing β -1.3-1.6glucan (*Aureobasidium* medium).

Fig. 3 shows the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the proliferation of U937 leukemia cells.

Fig. 4 shows the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the proliferation of Molt-4 leukemia cells.

Fig. 5 shows the influence of β -1.3-1.6glucan on the proliferation of P30/OHK leukemia cells.

Fig. 6 shows the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the proliferation of Raji leukemia cells.

Fig. 7, panels (a) and (b) show the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on cell surface molecules CD11a and CD49d, respectively.

Fig. 8, panels (a) and (b) show palm stamps of untreated hands and hands wiped with β -1.3-1.6glucan (*Aureobasidium* medium), respectively. Black marks depict the growth of microorganisms.

DETAILED DESCRIPTION OF THE INVENTION

Aureobasidium medium that contains β -1.3- 1.6glucan may be produced by strains of *Aureobasidium*. Advantageously, strain FERM P-18099 may be used. This strain has been deposited at the:

Agency of Industrial Science and Technology Life Engineering Research Institute,
Higashi 1-chome, 1-3, Tsukuba-shi,
Ibaraki, 305-0046, Japan.

The characteristics of this strain are described below.

5 Properties, taxonomy and mycological features of FERM P-18099. FERM P-18099
is a black fungus used for preparing *Aureobasidium* medium that contains β -1.3- 1.6glucan as
a major component. Its properties are as follows:

10 Scientific properties. This fungus produces high-molecular polysaccharide with high
viscosity. This substance agglutinates easily with ethanol, making it possible to collect
simply. Other purification methods, such as those described by U.S. Patent 5,789,579 for
purification of an α -type acid polysaccharide, may also be used. This polysaccharide is of β
type, and is acidic polysaccharide having a main chain of 1,3 bond and branches from 3- and
6-positions. It contains carboxylic acids such as malic acid as organic acids and phosphoric
acid. Moreover, it agglutinates easily with aluminum ions etc. This substance is also
effective for the promotion of growth as a feed and the effluent treatment. It is effective as a
food additive and functional food with immunity.

15 Taxonomic position. FERM P-18099 forms blackish brown colonies on
potato-dextrose-agar slant culturing for 7 days at 25 C. The fringe of colonies shows
filamentous growth and becomes gradually light blackish brown. The cells are filamentous,
20 and sometimes arthrospores, yeast-like budding conidiospores, oval yeast-like single cells,
and, in some time, thick-walled spore cells are formed. The growth temperature is 25°C, and
it decomposes hexoses such as glucose, fructose and galactose, sucrose, and starch. The
medium becomes conspicuously viscous. Based on FERM P-18099's mycological properties,
it is a kind of *Aureobasidium pullulans* in the black fungus family of deuteromycetes.

25 Mycological features of the isolated fungus. A colony of FERM P-18099 has a
smooth surface at first and grows into a grayish white, mucous and glossy oil drop-like
(fat-like), yeast-like material. The filamentous fungus body grows radially from the fringe
thereof, leading to crinkled, filamentous and just dendritic growth. This filamentous fungus
body grows well not only on the surface of medium, but also in the medium. In a short time,
30 light dark brown specks appear here and there on the surface of colony, which become black

specks gradually, and overall surface becomes dark black eventually. On this filamentous fungus body, a lot of light brown, elliptic or oval conidiospores are produced laterally. This conidiospore falls easily in pieces. While the surface of oil drop-like colony puts on the conidiospores here and there.

- 5 A medium containing sugars becomes very viscous and produces leather-like black culture moss on the surface of medium with thick colony formation. The optimum growth temperature is 20 to 25° C, and it yields alcohols and organic acids from sugars such as grape sugar and cane sugar. It gives out peculiar fragrance.

Cultural features. *Aureobasidium* may be cultured on solid or liquid media.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
10195
10200
10205
10210
10215
10220
10225
102

Morphological features. Young cells are transparent, filamentous, crinkled and dendritic. Black oval spore-like fungus bodies (filamentous) are produced laterally in places. Adherent black spore-like oil drop-like colonies are found. These structures are frangible and fall into pieces when given a shock.

5 Physiological features. The optimum growth temperature is 20 to 25°C. The culture yields viscous material from carbon sources such as, glucose, sucrose, etc. and alcohols and organic acids from sugars such as glucose. It gives out peculiar fragrance.

Method for making β -1.3-1.6glucan (*Aureobasidium* medium)

10 Based on the above information, the β -1.3-1.6glucan (*Aureobasidium* medium) to be used in the invention is β -glucan produced by the latest cultural technology, to which we added additional developments and improvements. *Aureobasidium* medium containing β -1.3-1.6glucan may be produced by culturing *Aureobasidium* in a solid or preferably a liquid culture medium supporting its growth. Such culture media include a carbon source, such as a sugar or carbohydrate, a nitrogen source, such as amino acids or a peptide mixture, and other nutrients, such as salts of sodium, potassium, phosphorous, iron, manganese, magnesium, calcium and sulfur. As mentioned above, culture media include potato-glucose medium and Czapek medium. Other culture conditions, such as those described in U.S. Patent 5,789,579 for culture of a strain producing an α -type acidic polysaccharide, may also be used.

20 A liquid culture is preferably agitated or aerated. The pH of the culture may range from 2 to 10 and the culture temperature preferably ranges from about 20 to 25° C. *Aureobasidium* medium containing β -1.3-1.6glucan may be harvested after 1 to 10 days, preferably at the point of maximal production of β -1.3-1.6glucan. Medium may be separated from cells by centrifugation, for instance, by an initial low speed centrifugation at 5,000 rpm and a high-speed centrifugation at 12,000 rpm. Preferably, the β -1.3-1.6glucan level in *Aureobasidium* medium is at least 10 times higher than B-glucan levels found in mushrooms. Culture medium may be used directly, after sterilization or filtration, or after further purification.

30 β -1.3-1.6glucan in β -1.3-1.6glucan in *Aureobasidium* medium may be further purified by agglutination with an organic solvent, such as ethanol, or by agglutination with metal ions,

such as aluminum ions. Of course, other convention purification methods may used to further purify the β -1.3-1.6glucan in *Aureobasidium* medium, including precipitation, filtration, chromatography and electrophoresis. Preferably, the β -1.3-1.6glucan is sterilized by convention methods, for instance, by heat treatment or filtration, prior to use.

5 While β -glucans may also be extracted from the cell walls of *Aureobasidium*, such cell wall extracts may lose their functional properties upon extraction and fractionation.

10 The β -1.3-1.6glucan (*Aureobasidium* medium) of the present invention exhibit unique physical and physiological properties different from other β -glucans produced by other types of fungi, making it particularly useful in commercial and industrial applications, especially in the fields of medical care, health and welfare and in the food industry. The present inventors have found that β -1.3-1.6glucan (*Aureobasidium* medium) produced with black yeast (*Aureobasidium*) exerts particular regulatory and modulatory effects on the immune system. For instance, β -1.3-1.6glucan (*Aureobasidium* medium) may be used to increase the numbers of lymphocytes. It may also be used to promote DNA synthesis (cell division), change the morphology of monocytic cells, and modify functional molecules, such as receptors, on the surface of cell membrane. β -1.3-1.6glucan (*Aureobasidium* medium) also reduces the proliferation of cancer cells and thus exerts an anticancer effect. Moreover, it has also been found that the β - 1.3-1.6glucan (*Aureobasidium* medium) inhibits the proliferation of microbes, such as microbes which adhere to skin surfaces like the hands and feet. Thus, it exerts an antimicrobial and infection-preventative effect.

20 β -1.3-1.6glucan (*Aureobasidium* medium) may be used in method for treating inflammatory, infectious or neoplastic diseases. These disease include, for instance, chronic articular rheumatism, collagen disease, newly-risen autoimmune disease, virus or bacterial infection, cancer, invasive or metastatic cancer, oncogenic virus, neuropathy, dementia, Alzheimer's disease, hepatic disease, cirrhosis, viral hepatic disease, cryptogenic intractable disease, life habit disease, diabetes, hypertension, arterial sclerosis, osteoporosis, hemorrhoids, or radiation damage. β -1.3-1.6glucan (*Aureobasidium* medium) may be administered by topically, orally, by application to a mucous membrane, by inhalation, or by parenteral, subcutaneous or intravenous administration. β -1.3-1.6glucan (*Aureobasidium* medium) may be formulated into an appropriate form, such as into a sterile or lyophilized form, for such modes of administration or into unit dosages. It may also optionally be

25
30

combined with other pharmaceutical products or drugs.

β -1.3-1.6glucan (*Aureobasidium* medium) may be used in food products.

Advantageously, the beneficial immunological and antimicrobial characteristics of this product may be useful for addition to foods for debilitated or immunocompromised patients, such as aseptic or germless foods for debilitated or immunocompromised patients, such as transplantation patients. For instance, it may be used in foods administered to patients in sterilized rooms after bone marrow transplantation or for burn victims. It may also be used in health foods or nutritional supplements for health maintenance and promotion. It may be used in foods for the aged or debilitated, such as patients who need to maintain their physical strength and disease resistance, or for persons with digestion or assimilation problems, such as deglutition disorders. It may also be used during other types of rehabilitation procedures, or for pregnant women or women under climacteric disturbance (such as menopause).

β -1.3-1.6glucan (*Aureobasidium* medium) may also be incorporated into health foods to maintain a subject's physiological or immunological vitality and balance.

The type of food product into which β -1.3-1.6glucan (*Aureobasidium* medium) may be applied or incorporated is not particularly limited and may include liquid, semisolid or solid food products. For instance, the β -1.3-1.6glucan (*Aureobasidium* medium) may be added to hot or cold drinks, including soft drinks, milk products, soy or rice milk products, fruit or vegetable juices; fermented milk products, such as cheeses and yoghurt; ice cream or other frozen products, gelatin or agar products; canned or preserved products, such as jams or jellies, condiments, including catsup, mustards, sauces or gravies, and relishes; canned meats, fruits or vegetables; cereal products, such as rice or wheat products, including noodles, pastas, breads and crackers and cookies. The β -1.3-1.6glucan (*Aureobasidium* medium) may also be formulated into special nutritional foods, such as hospital foods or nutritional formulas, or as into a nutritional supplement, for instance, into tablet or capsule, optional in combination with other nutrients, such as vitamins or minerals. Such capsules, tablets or pills, advantageously contain between 1 mg and 1,000 mg of β -1.3-1.6glucan (*Aureobasidium* medium). Solutions of the β -1.3-1.6glucan (*Aureobasidium* medium) of the present invention may also be applied or sprayed on the surfaces of foods, such as fresh fruits and vegetables to prevent or inhibit microbial contamination or spoilage.

The β -1.3-1.6glucan (*Aureobasidium* medium) may also be incorporated into personal

care products, such as toothpastes or mouthwash, or into a cosmetic, such as shampoo, soap, powder, such as a body or face powder, a gel, a film, a masque, a cosmetic cream, or a lotion. Such compositions and products may also be used to treat topical skin disorders, infections, or burns, including sunburns or UV irradiation, or to protect the connective skin tissue.

5 Products, including pharmaceutical, food, personal care and cosmetic products, contain about 0.001mg/ gr to 100mg/gr of β -1.3-1.6glucan from *Aureobasidium* (culture medium) and a preferable amount of *Aureobasidium* culture medium to be used in such products and compositions is from 0.1 to 100 parts by weight%. *Aureobasidium* culture medium preferably contains from about 3 to 5 mg/ml β -1.3-1.6glucan. Accordingly, such
10 products may also preferably contain from about .003 to 5 mg/ml β -1.3-1.6glucan. Foods preferably contain about 5 to 100 parts by weight/100 parts total weight of *Aureobasidium* culture medium.

β -1.3-1.6glucan (*Aureobasidium* medium) may be formulated into a pharmaceutical product, such as a product for modulating the immune system, or for treating a microbial or inflammatory disease. The β -1.3-1.6glucan (*Aureobasidium* medium) may be ingested, applied topically or parenterally, or administered directly to the site or biological compartment containing cancer cells. It may also be incorporated into a diagnostic product, as a auxiliary reagent (e.g. a buffer, stabilizer or preservative), or as a active component for the diagnosis of diseases associated with altered levels of glucans.

20 Moreover, as shown below, β -1.3-1.6glucan (*Aureobasidium* medium) may be used in products and compositions for the prevention of nosocomial infection or removal of environmental pollutants, for instance, into agents that prevent the propagation of microbes. It may also be used as a sanitizer or disinfectant, for instance, for cleaning the bodies of persons who are aged or infirm and cannot bathe. The *Aureobasidium* culture medium may
25 also be used undiluted as a disinfectant. Additionally, it may be used to treat hospital personnel, patients, surfaces, equipment, supplies or instruments in order to prevent nosocomial infections. Moreover, it may be used to remove environmental pollutants, such as internal secretion detractors, environmental hormones, toxins, heavy metals, and dioxins. As a sanitizer or disinfectant, β -1.3-1.6glucan (*Aureobasidium* medium) may be added to
30 water or other conventional cleaning or disinfectant solutions, including detergent or soap solutions.

EXAMPLES

The following examples illustrate the effects of β -1.3-1.6glucan (*Aureobasidium* medium) on the immune system.

To test these effects, peripheral lymphocytes were conventionally separated from peripheral blood of normal and healthy subjects. Separated lymphocytes were divided into control groups, which were not treated with β -1.3-1.6glucan (*Aureobasidium* medium) and into experimental groups treated with β -1.3-1.6glucan (*Aureobasidium* medium).

Control and experimental groups were cultured in an appropriate culture medium (RPMI 1640 containing 10% fetal calf serum (FCS) and 20% human serum (HS)) for a fixed period to investigate the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the numbers of lymphocytes, cell division of lymphocytes, production level of IgG antibody and on the production of cytokines. Moreover, morphological change of the peripheral blood lymphocytes was observed using a phase microscope. Moreover, the effects of β -1.3-1.6glucan (*Aureobasidium* medium) treatment on lymphocytic cell surface molecules (receptors) was determined.

Additionally, the effects of β -1.3-1.6glucan (*Aureobasidium* medium) on different types of cancer cells, such as different leukemia cell lines, were determined by comparison of control cancer cells receiving no treatment with β -1.3-1.6glucan (*Aureobasidium* medium), with experimental samples of the same types of cancer cells treated with β -1.3-1.6glucan (*Aureobasidium* medium). The number of lymphocytes was observed and calculated under phase microscope by using hemocytometer. See Fig. 1.

The capacity of control or treated lymphocytes to synthesize DNA was determined by a convention method. Namely, it was measured by the uptake or incorporation of tritiated thymidine (3H-thymidine) as measured by scintillation counting. Uptake is represented by the number of counts per minute, cpm. See Tables 4 and 5.

The production level of antibody from the lymphocytes in human peripheral blood with β -1.3-1.6glucan (*Aureobasidium* medium), that is, the production level of IgG antibody from B cells being immunocytes was determined by using enzyme antibody technique. Namely, according to the usual method, the amount of antibody in actual medium was determined by using double antibody enzyme antibody, enzyme immunoassay (EIA)

technique. There are many methods for determining the amount of antibody, for example, method using plaque technique, Ouchterlony technique, fluorescent antibody technique, etc. However, the EIA technique used in the examples has very high sensitivity and allows to determine even with small amount of material, hence it is a method used routinely as an immunological examination technique. See Table I which shows effects of β -1.3-1.6glucan (*Aureobasidium* medium) on IgG production.

For investigating the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on lymphocytes, the production levels of particular cytokines, such as interleukin and interferon, was determined by an enzyme-antibody technique. The enzyme-antibody technique measures the amounts of serum and cytokines in medium. Commercial kits for this technique are available from various firms. The levels of cytokines, such as interleukin 1, 2, 4, 6, and interferon were determined using this method. Evaluation of the effects of β -1.3-1.6glucan (*Aureobasidium* medium) on particular aspects of the immune system was done by a determination of the amounts of and dynamics of cytokine production or release. These results are presented in Tables II and III.

For investigating the morphological changes of lymphocytes treated with β -1.3-1.6glucan (*Aureobasidium* medium), the lymphocytes cultured with β -1.3-1.6glucan (*Aureobasidium* medium) are observed under phase microscope. Following this, their morphological change, in particular, shape of cells and change in the ruffle structure of cell surface are observed and taken into photographs. See Fig. 2.

With respect to the investigation on the change in dynamics of lymphocyte or surface molecule (receptor) of model cell cultured with β -1.3-1.6glucan (*Aureobasidium* medium), the flow cytometry technique (fluorescence activated cell sorter: FACS) is used. This advantageous method is capable of analyzing the dynamics of cell surface antigen and receptor as an immunologically analyzing means. This determination is conducted by using monoclonal antibody that reacts specifically with cell surface and antibody labeled with fluorescence, and, actually, the fluorescence intensity is image analyzed with computer after the reflection of laser beam was taken-in, and the results are converted to histogram. See Fig. 7.

The influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the proliferation of

cancer cells, in particular, leukemia cells is judged by the life and death of cancer cells cultured with β -1.3-1.6glucan (*Aureobasidium* medium) for a fixed period. Namely, according to this method, the leukemia cells cultured with β -1.3-1.6glucan (*Aureobasidium* medium) are collected and the life and death of cancer cells are counted by using hemocytometer to investigate the extent of proliferation of cancer cells. In some experiments, cells are stained actually by using stain solution to investigate the influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the extent of proliferation of cancer cells. See Figs. 3-6.

For investigating the effect of β -1.3-1.6glucan (*Aureobasidium* medium) on the removal of environmental microbes, a palm stamp on a conventional agar medium was used. This method counts the number of microbial colonies on the palm stamp on the agar medium as a measure of number of environmental microbes adhered to the skin surface. Conventional tryptosoy agar medium (medium: trypton, soy peptone, sodium chloride, agar) was used as an agar medium. However, different media may be selected, for instance, for investigating staphylococcus (a main microbe in nosocomial infections), mannite- salt-agar medium is used. For *Salmonella*, coliform bacteria, etc. involved in food poisoning, a selective agar media such as SS (*Salmonella-Shigella*) agar medium and Drigalski agar medium, and the like, may be used. In respective agar media, pH indicators such as phenol red and BTB (bromthymol blue) are contained, making it possible to know the proliferation of microbes by judging the color of medium. See Fig. 8.

Example 1. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on numbers of lymphocytes in peripheral blood

Various concentrations of β -1.3-1.6glucan (*Aureobasidium* medium) were added to the lymphocytes in healthy and normal human peripheral blood, which were cultured for 6 to 7 days at 37°C. After culture, the number of lymphocytes was counted by using hemocytometer. As a result, as shown in Fig.1, β -1.3-1.6glucan (*Aureobasidium* medium) at a fixed concentration tended to increase the number of lymphocytes in peripheral blood about 1.2 to 1.5 times compared with control group (group without β -1.3-1.6glucan).

Moreover, lymphocytes in human peripheral blood were taken out from healthy and normal person who continued to drink β -1.3-1.6glucan (*Aureobasidium* medium) for longer

than 3 years and an optimum concentration of β -1.3-1.6glucan (*Aureobasidium* medium) was added again *in vitro* to investigate the morphological change thereof. Conspicuous blast formation (image of cell division) was recognized for the group with β -1.3-1.6glucan (*Aureobasidium* medium) added, compared with the group without addition (Fig.2: Photographs).

Example 2. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on production of IgG antibody

Various concentrations of β -1.3-1.6glucan (*Aureobasidium* medium) were added to the lymphocytes in healthy and normal human peripheral blood, which were cultured for 7 days at 37°C. After culture, the concentration of antibody (IgG) in the supernatant of medium was examined by enzyme antibody technique using the double antibody EIA technique. No significant effect on the production of human IgG antibody by lymphocytes in human peripheral blood was seen. This result suggests that β -1.3-1.6glucan (*Aureobasidium* medium) is unlikely to cause adverse effects, such as decreases in antibody production *in vivo*.

Table 1 Influence of β -1.3-1.6glucan (*Aureobasidium* medium) administration IgG antibody levels.

| Concentration of β -1.3-1.6glucan (mg/ml) | Concentration of IgG antibody produced (ng/ml) |
|---|--|
| 0 | 668 |
| 7.5 | 835 |
| 0.75 | 592 |
| 0.35 | 663 |
| 0.18 | 553 |

Example 3. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the production of cytokines IL-1 β or IL-2.

A fixed concentration (final concentration 400 times) of β -1.3-1.6glucan (*Aureobasidium* medium) was added to the lymphocytes in healthy and normal human peripheral blood, which was cultured for 5 days at 37°C. After culture, the concentration of cytokines (interleukin 1 : IL-1 and interleukin-2: IL-2) in the supernatant of medium was examined by enzyme antibody technique (enzyme immunoassay: EIA technique). As a result, as shown in Table 2, no remarkable difference was recognized in the level of production between the group with β -1.3-1.6 glucan (*Aureobasidium* medium) added and the group without addition. This result also suggests that the β -1.3-1.6 glucan (*Aureobasidium*) would be unlikely to cause adverse effects *in vivo*, such as effects associated with the production of IL-1 β or IL-2.

Table 2 Influence of β -1.3-1.6 glucan (*Aureobasidium* medium) on the production of cytokines

| Culture | Production of cytokine | |
|--|------------------------|-----------|
| | IL-1 β | IL-2 |
| No addition | <10 pg/ml | < 0.8U/ml |
| Addition with β -1.3-1.6 glucan (<i>Aureobasidium</i> medium) | <10 pg/ml | <0.8U/ml |

Example 4. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the production of cytokine (2)

A fixed concentration (final concentration 20 times) of β -1.3-1.6glucan (*Aureobasidium* medium) was added to the monocytic model cell strain, which was cultured for 2 to 4 days at 37°C. After culture, the concentration of various cytokines (interleukin 1 β : IL-1 β , interleukin-4: IL-4, interleukin-2: IL-2, interleukin-6: IL-6 and interferon γ : IFN- γ) in the supernatant of medium was examined by enzyme antibody technique (enzyme immunoassay: EIA technique). As a result, as shown in Table 3, about 20 times or higher level of production of IL-6 was recognized for the group with β -1.3-1.6glucan (*Aureobasidium* medium) added, compared with the group without addition. However, these results suggest that administration of β -1.3-1.6 glucan would not be expected to induce side-effects associated with interleukin 1 β (IL-1 β), interleukin-4 (IL-4), interleukin-2 (IL-2), or interferon γ (IFN- γ).

Table 3 Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the production of various cytokines

| Culture | Production of cytokine | | | | |
|---|------------------------|----------|-----------|-----------|---------------|
| | IL-1 β | IL-2 | IL-4 | IL-6 | IFN- γ |
| No addition | <10pg/ml | <0.8U/ml | <2.0pg/ml | 2.0pg/ml | <0.1IU/ml |
| Addition with β -1.3-1.6glucan (<i>Aureobasidium</i> medium) | <10pg/ml | <0.8U/ml | <2.0pg/ml | 49.5pg/ml | <0.1IU/ml |

Example 5. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on normal lymphocyte cell division as measured by DNA synthesis.

β -1.3-1.6 glucan (*Aureobasidium* medium) (final concentration 400 times) was added to lymphocytes from normal and healthy human peripheral blood and also to the lymphocytes of a patient after bone marrow transplantation. The lymphocyte samples were cultured for 5 days at 37°C. After 5 days, tritiated thymidine (1 μ Ci) was added to these samples.

As shown in Table 4, normal lymphocytes treated with β -1.3-1.6glucan (*Aureobasidium* medium) exhibited increased cell division as measured by an average stimulation of DNA synthesis of about 4.5-times, compared to untreated normal lymphocytes.

Table 5 shows that lymphocytes from a majority of bone marrow transplantation patients also showed stimulation of cell division as measured by DNA synthesis. The stimulation index in Tables 4 and 5 represents the ratio of counts (cpm) in lymphocyte samples that were treated with β -1.3-1.6glucan (*Aureobasidium* medium) over untreated controls.

Table 4 Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on cell division as measured by DNA synthesis in lymphocytes from healthy and normal subjects.

| Lymphocyte in peripheral blood | Capability of DNA synthesis (cpm) | | SI (Stimulation index) |
|--------------------------------|-----------------------------------|--|---------------------------|
| | No addition | Addition with β -1.3-1.6glucan (<i>Aureobasidium</i> medium) | |
| SA | 624 | 851 | 1.4 |
| TS | 444 | 2067 | 4.7 |
| YA | 460 | 1232 | 2.7 |
| IS | 238 | 2177 | 9.1 |

cpm: Count per minute

SI: Stimulation Index

Table 5 Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on cell division as measured by DNA synthesis in patients after bone marrow transplantation.

| Lymphocyte | Transplantation form | Capability of DNA synthesis stimulation index (SI) | |
|------------|--|---|-----------------------|
| | | Stimulation with β -1.3-1.6glucan (<i>Aureobasidium</i> medium) | |
| | | Primary Stimulation | Secondary Stimulation |
| SU | Patient with bone marrow transplantation (Allogenic bone marrow transplantation) | 2.7 | 2.7 |
| AI | Patient with bone marrow transplantation (Allogenic bone marrow transplantation) | 3.6 | 5.2 |
| KM | Patient with bone marrow transplantation (Allogenic bone marrow transplantation) | 3.6 | 2.1 |
| NO | Patient with bone marrow transplantation (Autologous bone marrow transplantation) | 2.3 | 2.7 |
| TM | Patient with bone marrow transplantation (Allogenic bone marrow transplantation) | 1.2 | 0.6 |
| ZT | Patient with bone marrow transplantation (Autologous bone marrow transplantation) | 2.0 | 2.2 |
| YK | Patient with cord marrow transplantation (Allogenic transplantation) | 1.0 | 1.3 |
| SF | Patient with cord marrow transplantation (Allogenic transplantation) | 1.2 | 0.9 |

Example 6. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the proliferation of cancer cells

5 β -1.3-1.6Glucan (*Aureobasidium* medium) (final concentration 20 times) was added to various leukemia cells, which were cultured for 1 to 3 days at 37°C. After culture, the number of leukemia cells was counted by using hemocytometer to investigate the influence on the proliferation of leukemia cell. As shown in Figs 3-6, addition of β -1.3-1.6glucan (*Aureobasidium* medium) directly inhibited the proliferation of cancer (leukemia) cells. These data suggest the direct killing of cancer (leukemia) cells, for instance, by induction of
10 programmed death (apoptosis), as other components of the immune system were absent.

Example 7. Influence of β -1.3-1.6glucan (*Aureobasidium* medium) on the dynamics of cell surface molecule (receptor molecule) in monocyte/macrophage model system

15 β -1.3-1.6Glucan (*Aureobasidium* medium) (final concentration 20 times) was added to the cells in a monocyte/macrophage model system. Cells were cultured for 24 hours at 37°C. After 24 hours, the cells were collected and the dynamics of various human cell membrane antigens (receptor: CD antigen) were analyzed by indirect fluorescence antibody technique (flow cytophotometry technique) that uses various monoclonal antibodies. As shown in Fig.7, the levels of CD11a and CD49d molecules on the surface of cells in monocyte/macrophage model system cultured with β -1.3-1.6glucan (*Aureobasidium*
20 medium) were down-regulated.

Example 8. Effect of β -1.3-1.6glucan (*Aureobasidium* medium) on the prevention of propagation (proliferation) of environmental microbes

25 The removal or inactivation of environmental microbes adhered to surfaces or organisms is an important subject in the fields of medical care, health and welfare. Such removal or inactivation is particularly important for the prevention of nosocomial infections. Therefore, the antimicrobial effects of β -1.3-1.6glucan (*Aureobasidium* medium) on the proliferation of environmental microbes adhered to hands, feet and body, was evaluated using a palm stamp technique involving contacting the palm of the hand with a solid microbial medium. The hands of control subjects were contacted to the medium for about 20 seconds.

The hands of those in the experimental group were first wiped with a solution containing β -1.3-1.6glucan (*Aureobasidium* medium) and dried for about 5 minutes prior to contact with the medium. Similarly to the control group, the hands were then contacted with the nutritional solid medium for about 20 seconds.

5 The solid medium was cultured for 1 to 2 days at room temperature, and the proliferation of microbes (number of colonies) was observed. As shown in Fig.8, the palm stamps of hands treated with β -1.3-1.6glucan (*Aureobasidium* medium) showed far fewer microbial contaminants. This shows that the β -1.3-1.6glucan (*Aureobasidium* medium) inhibits the proliferation of environmental microbes and can be used as a sanitizer or the
10 bodies of persons who are aged or infirm and cannot bathe and also for the prevention of nosocomial infections and removal of environmental pollutants.

Modifications and other embodiments

Various modifications and variations of the described glucan products, compositions and methods as well as the concept of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has
15 been described in connection with specific preferred embodiments, it should be understood that the invention as claimed is not intended to be limited to such specific embodiments. Various modifications of the described modes for carrying out the invention which are obvious to those skilled in the chemical, biological, medical, environmental, cosmetic or food
20 arts or related fields are intended to be within the scope of the following claims.

Incorporation by Reference

Each document, patent application or patent publication cited by or referred to in this disclosure is incorporated by reference in its entirety. Any patent document to which this application claims priority is also incorporated by reference in its entirety. Specifically,
25 priority documents Japanese Patent Application 2000-342310, filed November 9, 2000, is hereby incorporated by reference.